

DIVISION OF MICROBIOLOGY AND INFECTIOUS DISEASES (DMID) PEDIATRIC TOXICITY TABLES NOVEMBER 2007 DRAFT

Note: The following toxicity table is a DRAFT and designed to provide general guidance on parameters for monitoring safety in clinical trials. This toxicity table is not comprehensive and should not be applied directly to all trials.

When selecting a toxicity table, the following are some of the items that must be taken into consideration:

- The population being studied
 - Does the clinical trial evaluate healthy subjects, subjects with a particular disease or condition?
- The stage of test article development
 - Is the clinical trial a Phase I (is it for the first time in human subjects?) , II, III or IV?
- The type of test article
 - Does the clinical trial evaluate a drug, device, vaccine or other biologic agent?
- The prior human and preclinical experience with the test article
 - Are there any specific findings that require adjustment of the toxicity table?
 - Has it been approved for this indication in adult population?

Single site clinical trials evaluating healthy subjects should conform to the laboratory normal values at the single site. Multi-center clinical trials should reconcile among their laboratory normal values when evaluating a healthy volunteer population.

Please confer with the DMID protocol team and DMID's Office of Clinical Research Affairs when selecting or developing a toxicity table for a DMID-sponsored trial.

DIVISION OF MICROBIOLOGY AND INFECTIOUS DISEASES (DMID) PEDIATRIC TOXICITY TABLES

NOVEMBER 2007

DRAFT

ABBREVIATIONS: Abbreviations utilized in the Table:

ULN = Upper Limit of Normal	LLN = Lower Limit of Normal
R _x = Therapy	Req = Required
Mod = Moderate	IV = Intravenous
ADL = Activities of Daily Living	Dec = Decreased

ESTIMATING SEVERITY GRADE

For abnormalities NOT found elsewhere in the Toxicity Tables use the scale below to estimate grade of severity:

GRADE 1	Mild	Transient or mild discomfort (< 48 hours); no medical intervention/therapy required
GRADE 2	Moderate	Mild to moderate limitation in activity - some assistance may be needed; no or minimal medical intervention/therapy required
GRADE 3	Severe	Marked limitation in activity, some assistance usually required; medical intervention/therapy required, hospitalizations possible
GRADE 4	Life-threatening	Extreme limitation in activity, significant assistance required; significant medical intervention/therapy required, hospitalization or hospice care probable
GRADE 5	Death	

SERIOUS OR LIFE-THREATENING AEs

ANY clinical event deemed by the clinician to be serious or life-threatening should be considered a grade 4 event. Clinical events considered to be serious or life-threatening include, but are not limited to: seizures, coma, tetany, diabetic ketoacidosis, disseminated intravascular coagulation, diffuse petechiae, paralysis, acute psychosis, severe depression.

COMMENTS REGARDING THE USE OF THESE TABLES

- Standardized and commonly used toxicity tables (Division of AIDS, NCI's Common Toxicity Criteria (CTC), and World Health Organization (WHO)) have been adapted for use by the Division of Microbiology and Infectious Diseases (DMID) and modified to better meet the needs of participants in DMID trials.
- For parameters not included in the following Toxicity Tables, sites should refer to the "Guide For Estimating Severity Grade" located above.
- Criteria are generally grouped by body system.
- Some protocols may have additional protocol specific grading criteria, which will supercede the use of these tables for specified criteria.

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT**

**(Selected Values for children less than or equal
to 3 months of age – does not apply for preterm infants)**

For all parameters not listed on this table, please refer
to the DMID Toxicity Table for children > 3 months of age.

HEMATOLOGY				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Hemoglobin				
1-7 days old	13.0-14.0 g m/dL	12.0-12.9 g m/dL	<12 g m/dL	Cardiac Failure secondary to Anemia
8-21 days old	12.0-13.0 g m/dL	10.0-11.9 g m/dL	<10.0 g m/dL	Cardiac Failure secondary to Anemia
22-35 days old	9.5-10.5 g m/dL	8.0-9.4 g m/dL	<8.0 g m/dL	Cardiac Failure secondary to Anemia
36-60 days old	8.5-9.4 g m/dL	7.0-8.4 g m/dL	<7.0 g m/dL	Cardiac Failure secondary to Anemia
61-90 days old	9.0-9.9 g m/dL	7.0-8.9 g m/dL	<7.0 g m/dL	Cardiac Failure secondary to Anemia
Abs Neutrophil Ct				
1 day old	5000-7000/mm ³	3000-4999/mm ³	1500-2999/mm ³	<1500/mm ³
2-6 days old	1750-2500/mm ³	1250-1749/mm ³	750-1249/mm ³	<750/mm ³
7-60 days old	1200-1800/mm ³	900-1199/mm ³	500-899/mm ³	<500/mm ³
61-90 days old	750-1200/mm ³	400-749/mm ³	250-399/mm ³	<250/mm ³
Bilirubin (Fractionated bilirubin test must be performed when total bilirubin is elevated)				
<7 days old	.	20-25mg/dL	26-30 mg/dL	>30 mg/dL
7-60 days old	1.1-1.9xN	2.0-2.9Xn	3.0-7.5xN	>7.5xN
61-90 days old	1.1-1.9xN	2.0-2.9xN	3.0-7.5xN	>7.5xN

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT**

**(Selected Values for children less than or equal
to 3 months of age)**

HEMATOLOGY (continued)				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Creatinine				
<7 days old	1.0-1.7 mg/dL	1.8-2.4 mg/dL	2.5-3.0 mg/dL	>3.0 mg/dL
7-60 days old	0.5-0.9 mg/dL	1.0-1.4 mg/dL	1.5-2.0 mg/dL	>2.0 mg/dL
61-90 days old	0.6-0.8 mg/dL	0.9-1.1 mg/dL	1.2-1.5 mg/dL	>1.5 mg/dL
Cr Clearance				
<7 days old	35-40 ml/min	30-34 ml/min	25-29 ml/min	<25 ml/min
7-60 days old	45-50 ml/min	40-44 ml/min	35-39 ml/min	<35 ml/min
61-90 days old	60-75 ml/min	50-59 ml/min	35-49 ml/min	<35 ml/min
Hypocalcemia				
<7 days old	6.5-6.9 mEq/L	6.0-6.4 mEq/L	5.5-5.9 mEq/L	<5.5 mEq/L
7-60 days old	7.6-8.0 mEq/L	7.0-7.5 mEq/L	6.0-6.9 mEq/L	<6.0 mEq/L
61-90 days old	7.8-8.4 mEq/L	7.0-7.7 mEq/L	6.0-6.9 mEq/L	<6.0 mEq/L
Hypercalcemia				
<7 days old	12.0-12.4 mEq/L	12.5-12.9 mEq/L	13.0-13.5 mEq/L	>13.5 mEq/L
7-60 days old	10.5-11.2 mEq/L	11.3-11.9 mEq/L	12.0-13.0 mEq/L	>13.0 mEq/L
61-90 days old	10.5-11.2 mEq/L	11.3-11.9 mEq/L	12.0-13.0 mEq/L	>13.0 mEq/L

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT**

(Greater than 3 months of age)

LOCAL REACTIONS				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Induration	< 10mm	10-25 mm	26-50mm	>50mm
Erythema	< 10mm	10-25 mm	26-50mm	>50mm
Edema	< 10mm	10-25 mm	26-50mm	>50mm
Rash at Injection Site	< 10mm	10-25 mm	26-50mm	>50mm
Pruritus	Slight itching at injection site	Moderate itching at injection extremity	Itching at injection extremity and other sites	Itching over entire body

HEMATOLOGY				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Hemoglobin for children greater than months and less than 2 years of age	9.0-9.9 gm/dL	7.0-8.9 gm/dL	<7.0 gm/dL	Cardiac Failure secondary to anemia
Hemoglobin for children greater than 2 years of age	10-10.9 gm/dL	7.0-9.9 gm/dL	<7.0 gm/dL	Cardiac Failure secondary to anemia
Absolute Neutrophil Count	750-1200/mm ³	400-749/mm ³	250-399/mm ³	<250/mm ³
Platelets	-----	50,000-75,000/mm ³	25,000-49,999/mm ³	<25,000/mm ³
Prothrombin Time (PT)	1.1-1.2 x ULN	1.3 -1.5 x ULN	1.6 -3.0 x ULN	>3.0 x ULN
Partial Thromboplastin Time (PTT)	1.1-1.6 x ULN	1.7-2.3 x ULN	2.4 -3.0 x ULN	>3.0 x ULN

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT
(Greater than 3 months of age)**

GASTROINTESTINAL				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Bilirubin (when accompanied by any increase in other liver function test)	1.1 - <1.25 x ULN	1.25 - <1.5 x ULN	1.5 – 1.75 x ULN	> 1.75 x ULN
Bilirubin (when other liver function are in the normal range)	1.1 - <1.5 x ULN	1.5 - <2.0 x ULN	2.0 – 3.0 x ULN	> 3.0 x ULN
AST (SGOT)	1.1 - <2.0 x ULN	2.0 – <3.0 x ULN	3.0 – 8.0 x ULN	> 8 x ULN
ALT (SGPT)	1.1 - <2.0 x ULN	2.0 – <3.0 x ULN	3.0 – 8.0 x ULN	> 8 x ULN
GGT	1.1 - <2.0 x ULN	2.0 – <3.0 x ULN	3.0 – 8.0 x ULN	> 8 x ULN
Pancreatic Amylase	1.1-1.4 x ULN	1.5-1.9 x ULN	2.0-3.0 x ULN	>3.0 x ULN
Uric Acid	7.5-9.9mg/dL	10-12.4 mg/dL	12.5-15.0 mg/dL	>15.0 mg/dL
CPK	See Neuro muscular Toxicity			
Appetite	-----	Decreased appetite	Appetite very decreased, no solid food taken	No solid or liquid taken
Abdominal Pain	Mild	Moderate- No Treatment Needed	Moderate- Treatment Needed	Severe- Hospitalized for treatment
Diarrhea	Slight change in consistency and/or frequency of stools	Liquid stools	Liquid stools greater than 4x the amount or number normal for this child	Liquid stools greater than 8x the amount or number normal for this child

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT
(Greater than 3 months of age)**

GASTROINTESTINAL (continued)				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Constipation	Slight change in the consistency/frequency of stool	Hard, dry stools with a change in frequency	Abdominal pain	Distention and Vomiting
Nausea	Mild	Moderate- Decreased oral intake	Severe-Little oral intake	Unable to ingest food or fluid for more than 24 hours
Vomiting	1 episode/day	2-3 episodes per day	4-6 episodes per day	Greater than 6 episodes per day or Intractable Vomiting

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT
(Greater than 3 months of age)**

ELECTROLYTES				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
CREATININE				
3 Months -2 Years of age	0.6-0.8 x ULN	0.9-1.1 x ULN	1.2-1.5 x ULN	>1.5 x ULN
2 Years- 12 Years of age	0.7-1.0 x ULN	1.1-1.6 x ULN	1.7-2.0 x ULN	>2.0 x ULN
Greater than 12 Years of age	1.0-1.7 x ULN	1.8-2.4 x ULN	2.5-3.5 x ULN	>3.5 x ULN

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT**

(Greater than 3 months of age)

ELECTROLYTES				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Hypernatremia		<145-149 mEq/L	150-155 mEq/L	>155 mEq/L or abnormal sodium AND mental status changes
Hyponatremia		130-135 mEq/L	129-124 mEq/L	<124 mEq/L or abnormal sodium AND mental status changes
Hyperkalemia	5.0-5.9 mEq/L	6.0-6.4 mEq/L	6.5-7.0 mEq/L	>7.0 mEq/L or abnormal potassium AND cardiac arrhythmia
Hypokalemia	3.0-3.5 mEq/L	2.5-2.9 mEq/L	2.0-2.4 mEq/L	<2.0 mEq/L or abnormal potassium AND cardiac arrhythmia
Hypercalcemia	10.5-11.2mg/dL	11.3-11.9 mg/dL	12.0-12.9 mg/dL	>13.0 mg/dL
Hypocalcemia	7.8-8.4 mg/dL	7.0-7.7 mg/dL	6.0-6.9 mg/dL	<6.0 mg/dL
Hypomagnesemia	1.2-1.4 mEq/L	0.9-1.1 mEq/L	0.6-0.8 mEq/L	<0.6 mEq/L or abnormal magnesium AND cardiac arrhythmia
Hypoglycemia	55-65 mg/dL	40-54 mg/dL	30-39 mg/dL	<30 mg/dL or abnormal glucose AND mental status changes
Hyperglycemia	116-159 mg/dL	160-249 mg/dL	250-400 mg/dL	>400 mg/dL or ketoacidosis
Proteinuria	Tr- 1+ or <150 mg/day	2+ or 150-499 mg/day	3+ or 500-1000 mg/day	4+ or Nephrotic syndrome >1000 mg/day
Hematuria	Microscopic <25	Microscopic >25		Gross hematuria

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT**

	cells/hpf	cells/hpf		
--	-----------	-----------	--	--

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT
(Greater than 3 months of age)**

CENTRAL NERVOUS SYSTEM (CNS)				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Generalized CNS Symptoms			Dizziness	Hypotonic, hyporesponsive episodes; Seizures; Apnea/Bradycardia; Inconsolable crying > 3 hrs;
Headache	Mild	Moderate, Responds to non-narcotic analgesia	Moderate to Severe, Responds to narcotic analgesia	Intractable
Level of Activity		Slightly irritable OR slightly subdued	Very irritable OR Lethargic	Inconsolable OR Obtunded
Visual		Blurriness, diplopia, or horizontal nystagmus of < 1 hour duration, with spontaneous resolution	More than 1 episode of Grade 2 symptoms per week, or an episode of Grade 2 symptoms lasting more than 1 hour with spontaneous resolution by 4 hours or vertical nystagmus	Decrease in visual acuity, visual field deficit, or oculogyric crisis
Myelopathy		None	None	Myelopathic/spinal cord symptoms, such as: pyramidal tract weakness and disinhibition, sensory level, loss of proprioception, bladder/bowel dysfunction

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT
(Greater than 3 months of age)**

PERIPHERAL NERVOUS SYSTEM				
PARAMETER	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Neuropathy/ Lower Motor Neuropathy		Mild transient Paresthesia only	Persistent or progressive paresthesias, burning sensation in feet, or mild dysesthesia; no weakness; mild to moderate deep tendon reflex changes; no sensory loss	Onset of significant weakness, decrease or loss of DTRs, sensory loss in "stocking glove" distribution, radicular sensory loss, multiple cranial nerve involvement; bladder or bowel dysfunction, fasciculations, respiratory embarrassment from chest wall weakness.
Myopathy or Neuromuscular Junction Impairment	Normal or mild ($<2 \times$ ULN) CPK elevation	Mild proximal weakness and/or atrophy not affecting gross motor function. Mild myalgias, +/- mild CPK elevation ($<2 \times$ ULN)	Proximal muscle weakness and/or atrophy affecting motor function +/- CPK elevation; or severe myalgias with CPK $>2 \times$ ULN;	Onset of myasthenia- like symptoms (fatigable weakness with external, variable ophthalmoplegia and/or ptosis), or neuromuscular junction blockade (acute paralysis) symptoms

**DIVISION OF MICROBIOLOGY AND INFECTIOUS
DISEASES (DMID) PEDIATRIC TOXICITY TABLES
NOVEMBER 2007
DRAFT
(Greater than 3 months of age)**

OTHER				
	GRADE 1	GRADE 2	GRADE 3	GRADE 4
Allergy	Pruritus without Rash	Pruritic Rash	Mild Urticaria	Severe Urticaria Anaphylaxis, Angioedema
Drug Fever (Rectal)	.	38.5-40C 101.3 – 104.0F	Greater than 40.0C Greater than 104.0F	Sustained Fever: Equal or greater than 40C (104.0F) for longer than 5 days
Cutaneous	Localized rash	Diffuse macu lopapular Rash	Generalized urticaria	Stevens-Johnson Syndrome or Erythe ma multi forme
Stomatitis	Mild discomfort	Painful, difficulty swallowing, but able to eat and drink	Painful: unable to swallow solids	Painfu l: unable to swallow liquids; requires IV fluids
Clinical symptoms <i>not otherwise specified</i> in this table	No therapy; monitor condition	May require minimal intervention and monitoring	Requires medical care and possible hospitalization	Requires active medical intervention, hospitalization, or hospice care
Laboratory values <i>not otherwise specified</i> in this table	Abnormal, but requiring no immediate intervention; follow	Sufficiently abnormal to require evaluation as to causality and perhaps mild therapeutic intervention, but not of sufficient severity to warrant immediate changes in study drug	Sufficiently severe to require evaluation and treatment, including at least temporary suspension of study drug	Life-threatening severity; Requires immediate evaluation, treatment, and usually hospitalization; Study drug must be stopped immediately and should not be restarted until the abnormality is clearly felt to be caused by some other mechanis m that study drug